

Invariants of tangles with flat connections in their complements.I. Invariants and holonomy R -matrices

R. Kashaev, N.Reshetikhin

February 1, 2008

Abstract

The notion of holonomy R -matrices is introduced. It is shown how to define invariants of tangles with flat connections in a principle G -bundle of the complement of a tangle using holonomy R -matrices.

1 Introduction

A breakthrough in the theory of invariants of knots came with the discovery of the Jones polynomial [J] and its generalizations (HOMFLY and Kauffman). The construction of the Jones invariant based on a solution to the Yang-Baxter equation was given in [J-1] and was generalized to HOMFLY polynomial and to more general case in [T]. It involved a solution to the Yang-Baxter equation which satisfies certain extra condition. Then it was shown in [R] [RT] that such invariants can be obtained from quantized universal algebras for any simple Lie algebra. In [R-1] it was proven that the non-degeneracy and the cross-nondegeneracy of a solution of the Yang-Baxter equation is enough to construct the invariants.

In this paper we re-examine the construction of invariants of links from [RT] and [R-1] (for details see [T-1]. Using systems of holonomy R -matrices we construct the invariant of tangles in $\mathbb{R}^2 \times I$ (here I is a unit interval) with a flat connection in a principle G -bundle (where G is a simple complex algebraic group) over the complement of a tangle. We show that this invariant depends only on a gauge class of this connection.

In the follow-up paper we will show that quantized universal R -matrices provide examples of holonomy R -matrices.

In the second section of this paper we describe the moduli space of flat connections in the complement to a tangle in $\mathbb{R}^3$. In the third section we describe the category of tangles with flat connections in the complement. This construction is very similar to π -tangles

introduced and studied in [T-2]. Then in section 4 we construct invariants of tangles with flat connections in the complement using non-degenerate solutions to the holonomy Yang-Baxter equations. In section 5 we show that these invariants are gauge invariant, and in particular for links they can be functions (sections of a line bundle) over the moduli space of flat connections in the complement of tangle. Some proofs are moved to the Appendix.

This paper is one of a series of papers in which we will analyze invariants of tangles and 3-manifolds related to quantized universal enveloping algebras with large center. Some outline of this program is given in the conclusion (section 6).

The main results of this paper and of the part II were announced at the conference "Graphs and Patterns in Mathematics and Physics". Both authors were partly supported by the NSF grant DMS-0070931 and by the CRDF grant RM1-2244.

2 Flat connections in the complement of a tangle

2.1 The fundamental group of the complement of a tangle

Let $I \equiv [0, 1]$ be the unit closed interval. A geometric tangle $t \subset \mathbb{R}^2 \times I$ is the image of an embedding

$$\underbrace{I \sqcup \dots \sqcup I}_{k \text{ times}} \sqcup \underbrace{S^1 \sqcup \dots \sqcup S^1}_{l \text{ times}} \rightarrow \mathbb{R}^2 \times I$$

with oriented components such that

$$\partial t \subset \mathbb{R}^2 \times \partial I = \mathbb{R}^2 \times \{0, 1\}$$

and $\partial t \cap \mathbb{R}^2 \times \partial I = t \cap \mathbb{R}^2 \times \partial I$.

Let (x, y, z) , $x, y \in \mathbb{R}$, $z \in I$, be Cartesian coordinates on $\mathbb{R}^2 \times I$. Denote by D_t the image of t under the projection

$$p: \mathbb{R}^2 \times I \ni (x, y, z) \mapsto (y, z) \in \mathbb{R}^2 \quad (1)$$

We call geometric tangle t *restricted* if it is based on lines $L_{\pm} = \{(0, y, (1 \pm 1)/2)\}$, i.e. $\partial t \in L_+ \cup L_-$. Denote by

$$\partial_{\pm} t \equiv \partial t \cap L_{\pm}, \quad \partial_{\pm} D_t \equiv p(\partial_{\pm} t)$$

the boundary components of t and D_t which belong to $L_{\pm}$ and their images in D_t , respectively.

Projection D_t is called *regular* if it has only double points at which segments intersect transversely, all critical points in z -direction are non-degenerate, and non of the critical




Figure 1:




Figure 2:

points coincide with any of the double points. A regular projection D_t will be called *diagram* of t . The vertices of a diagram are its double points, critical points in z -direction and boundary points, while edges are the segments running between the vertices. A vertex of D_t is called *positive or (+) crossing* if it is a double point and the angle between lower and upper components is positive. If the angle is negative, the vertex is called *negative or (-) crossing*, see Fig. 1.

For each double point v of diagram D_t denote by a_v, b_v, c_v, d_v the adjacent edges in accordance with Fig. 2, and for each critical point v denote by l_v, r_v the adjacent edges as in Fig. 3. Define group $\Pi(D_t)$ generated by edges of D_t with the following defining relations:

- for each double point v

$$\begin{aligned}
 (+) \text{ crossing: } & a_v = d_v, \quad b_v = d_v^{-1} c_v d_v \\
 (-) \text{ crossing: } & b_v = c_v, \quad a_v = c_v d_v c_v^{-1}
 \end{aligned}$$




Figure 3:




Figure 4: The choice of the monodromy path

- for each critical point v : $l_v r_v = 1$.

The isomorphism class of this group does not depend on the diagram representing t and is isomorphic to the fundamental group of the complement to t :

$$\Pi(D_t) \simeq \pi_1(\mathbb{R}^2 \times I \setminus t)$$

In fact, the definition of $\Pi(D_t)$ concides with the Wirtinger presentation of $\pi_1(\mathbb{R}^2 \times I \setminus t)$. One can fix an isomorphism by choosing a base point x_0 in $\mathbb{R}^2 \times I \setminus t$ and associating to each edge e in D_t element $\Gamma_e^{x_0}$ of $\pi_1(\mathbb{R}^2 \times I \setminus t, x_0)$ which is represented by the loop at x_0 with projection as in Fig. 4. We shall call such correspondence a *monodromy* map.

Associated with boundary of D_t , group $\Pi(D_t)$ has two natural subgroups $\Pi(\partial_{\pm} D_t)$ generated by edges intersecting with corresponding boundary components. One has the following commutative diagram:

$$\begin{array}{ccccc} \Pi(\partial_- D_t) & \hookrightarrow & \Pi(D_t) & \hookleftarrow & \Pi(\partial_+ D_t) \\ \uparrow & & \wr & & \uparrow \\ \pi_1(\mathbb{R}^2 \times \{0\} \setminus \partial_- t, x_0^-) & \rightarrow & \pi_1(\mathbb{R}^2 \times I \setminus t, x_0) & \leftarrow & \pi_1(\mathbb{R}^2 \times \{1\} \setminus \partial_+ t, x_0^+) \end{array}$$

where the hookarrows are embeddings while the arrows are natural group homomorphisms, and we denote $x_0^\pm = (x, y, (1 \pm 1)/2)$ the projections of the base point $x_0 = (x, y, z)$. The base point is chosen such that it projects to the plane of the diagram D_t to the far left of it (in the y -direction).

2.2 The moduli space of flat connections in the complement to a tangle

For a Lie group G let A be the 1-form corresponding to a flat connection in a principle G -bundle over the complement to tangle t in $\mathbb{R}^2 \times I$. Such form defines a group homomorphism $\pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \rightarrow G$ which maps the element generated by path Γ to the holonomy of A along Γ . Because we have the isomorphism $\pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \simeq \Pi(D_t)$ a flat G -connection also defines a group homomorphism

$$\gamma: \Pi(D_t) \ni e \mapsto \gamma_e \in G$$

We denote the space of all such homomorphisms as $\Pi(D_t, G)$. The group G acts on $\Pi(D_t, G)$ by conjugations

$$g: \gamma_e \mapsto g\gamma_e g^{-1}$$

In terms of $\pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \simeq \Pi(D_t)$ this action corresponds to the changes of the trivialization of the G -bundle at the base point, i.e. to gauge transformations.

Thus, the gauge class of the flat connection A defines an element in the coset space $\Pi(D_t, G)/G$. This gives a combinatorial description of the space of isomorphism classes of representations of the fundamental group of $\mathbb{R}^2 \times I \setminus t$ in G (or equivalently, of the moduli space of flat connections):

$$\Pi(D_t, G)/G \simeq \{\pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \rightarrow G\}/G$$

Notice that this description of the isomorphism classes of representations of the fundamental group works also in cases when G is not a Lie group.

Notice that the homomorphism $f: \pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \rightarrow G$ does not define a flat connection in the complement of the tangle. It only defines a gauge class of a flat connection.

For what will follow we need slightly more then a representation of the fundamental group, but less then a flat connection. Let $\tilde{x}_0$ be a point which projects to the far right from D_t . A pair (f, g) where f is a representation of the fundamental group and $g \in G$ is called a *ramified representation of the fundamental group* if g has a meaning of the holonomy of




Figure 5:

a flat connection whose gauge class is defined by f along the path connection x_0 and $\tilde{x}_0$ in such a way that it projects "over" the diagram D_t (see Fig. 5).

Let $\{g_e\}_{e \in E(D_t)}$ be a ramified representation of the fundamental group. The gauge group acts on it as $g_e \mapsto gg_eg^{-1}$, $g_r \mapsto gg_rh^{-1}$. Thus the action of the gauge group on ramified representations reduces to the action of $G \times G$. The isomorphism class does not depend on the ramification.

2.3 Factorizable groups and Lie groups

We say that group G is factorizable into two subgroups $G_{\pm} \subset G$ if any element $g \in G$ can be represented in a unique way as

$$g = g_+g_-^{-1} = \bar{g}_-^{-1}\bar{g}_+ \quad (2)$$

where $g_{\pm}, \bar{g}_{\pm} \in G_{\pm}$.

Let G be a Lie group with Lie algebra $\mathfrak{g}$. We say that Lie algebra $\mathfrak{g}$ is factorizable into two Lie subalgebras $\mathfrak{g}_{\pm}$ if for each $x \in \mathfrak{g}$ there exists unique pair $x_{\pm} \in \mathfrak{g}_{\pm}$ such that

$$x = x_+ - x_-$$

A Lie group with factorizable Lie algebra is almost factorizable (i.e. factorizable on an open dense subset). It will be called factorizable Lie group. We will be especially interested in the case when G is an algebraic Lie group with factorizable Lie algebra. Then it is factorizable on a Zariski open subset of G .

Another important case is the conditional factorization. In conditionally factorizable Lie algebras factorization (2) is unique only under extra conditions on $x_{\pm}$. For example, fixing a Borel subalgebra $\mathfrak{b}_+$ in a simple Lie algebra $\mathfrak{g}$ makes $\mathfrak{g}$ conditionally factorizable with $\mathfrak{g}_{\pm} = \mathfrak{b}_{\pm}$ and with the condition $x_+|_{\mathfrak{h}} = -x_-|_{\mathfrak{h}}$, where $\mathfrak{h}$ is the corresponding Cartan subalgebra.

Remark 1. *We can choose $G_+ = \{e\}$ and $G_- = G$. We call it trivial factorizability.*

Let G be a factorizable group. Define a binary operation

$$g \star h = g_+ h_+ (g_- h_-)^{-1}$$

which obviously defines a group structure on G with the same identity element as for the original group structure. The inverse to g in this group is $i(g) = g_+^{-1} g_-$. For factorizable Lie groups this structure is defined on a dense open subset of G (or on $G_+ \times G_-$).

2.4 Combinatorial description of the moduli space for factorizable groups

Assume G is a factorizable group and D_t is a regular diagram of a tangle.

Definition 1. *The map $E(D_t) \rightarrow G$ which associates to edge e element $x_e \in G$ is called a G -coloring of diagram D_t if it satisfies the relations:*

- *double point:* $x_{b_v} = (x_{a_v})_{\pm}^{-1} x_{c_v} (x_{a_v})_{\pm}$, $x_{d_v} = (x_{c_v})_{\mp}^{-1} x_{a_v} (x_{c_v})_{\mp}$, for $(\pm)$ crossing,
- *critical point:* $x_{e_v} = x_{r_v}$.

Theorem 1. *There is one to one correspondence between G -colorings of a regular diagram of a tangle and generic representations of the fundamental group of the complement.*

Proof. We define a representation of $\pi_1(\mathbb{R}^2 \times I \setminus t)$ which maps the homotopy class of path Γ_e (see Fig. 4) to the element

$$(x_{e_1})_-^{\epsilon_1} \dots (x_{e_n})_-^{\epsilon_n} x_e (x_{e_n})_-^{-\epsilon_n} \dots (x_{e_1})_-^{-\epsilon_1} \quad (3)$$

where $e_1, \dots, e_n$ is the sequence of edges, intersected by corresponding to edge e monodromy path (enumeration is given along the opposite direction to its orientation), while ϵ_i are the signs defined by the projection of edges e_i to vertical axis:

$$\epsilon_i = \begin{cases} 1 & \text{if } e_i \text{ is oriented upwards} \\ -1 & \text{if } e_i \text{ is oriented downwards} \end{cases}$$




Figure 6: There horizontal lines are parts of the monodromy path while vertical lines are edges of the diagram of a tangle.

In other words, the holonomy along a monodromy path is computed as a product of elements corresponding to "elementary intersections" between edges of the diagram and this path, taken in the order opposite to the orientation of the path. Elements corresponding to the "elementary intersections" are given on Fig. 6. To prove the theorem one has to check that holonomies (3) satisfy the relations defining the set $\Pi(D_t, G)$ and that on an open dense subset of $\Pi(D_t, G)$ one can solve equations (3) recursively. These are simple straightforward calculations. $\square$

For factorizable G we will make the standard choice of ramification of a representation of the fundamental group of the complement. Assume that the representation is generic and that $\{x_e\}$ is the G -coloring corresponding to it. Then the standard ramification is obtained by applying rules from Fig. 6 to the path ramification path connection points x_0 and $\tilde{x}_0$:

$$g = (x_{e_1})_-^{-1} \dots (x_{e_n})_-^{-1}$$

Here $e_1, \dots, e_n$ are edges of D_t which intersect with the ramification path.

3 The category of tangles with flat connections in their complements

In this section we will define the category of tangles with ramified representations of the fundamental group of the complement. Since a flat connection defines a ramified representation of the fundamental group, by the abuse of language we will call this category the category of tangles with flat connections in the complement.

3.1 The equivalence

Restricted geometric tangle t is called *standard* if $\partial_{\pm} t = \{(0, 1, 1), (0, 2, 1) \dots (0, n_{\pm}, 1)\}$ for some $n_{\pm} \in \mathbb{Z}_{\geq 0}$. We identify planes $\mathbb{R}^2 \times \{0\}$ and $\mathbb{R}^2 \times \{1\}$ with $\mathbb{R}^2$ through the projection p defined in (1).

Consider the set of pairs (t, f, g) , where t is a standard tangle and $f : \pi_1(\mathbb{R}^2 \times I \setminus t, x_0) \rightarrow G$ is a group homomorphism and $g \in G$ is its ramification. Define the following equivalence relation on this set. Pair (t, f, g) is equivalent to (t', f', g) if t is isotopic to t' in $\mathbb{R}^2 \times I$ by an isotopy preserving the boundary and points $x_0, \tilde{x}_0$ (so, in particular, $\partial_+ t = \partial_+ t'$), which brings flat connection f to f' . Equivalence class $[t, f, g]$ will be called a (standard) tangle with a ramified representation of the fundamental group in the complement.

3.2 The composition

From now on the point x_0 will have coordinates $(0, y_0, 1/2)$ with $y_0 \ll -1$ and $\tilde{x}_0$ will have coordinates $(0, \tilde{y}_0, 1/2)$ with $\tilde{y}_0 \gg 1$.

Let t_1 and t_2 be two standard geometrical tangles with $\partial_+ t_1 = \partial_- t_2$. Define their composition $t_1 \circ t_2$ as the tangle with the following defining properties:

$$t_1 \circ t_2 \cap \mathbb{R}^2 \times [0, 1/2] \simeq t_1, \quad (4)$$

$$t_1 \circ t_2 \cap \mathbb{R}^2 \times [1/2, 1] \simeq t_2. \quad (5)$$

Here the identifications are given by linear maps, $(x, y, z) \rightarrow (x, y, 2z)$ for the first one, and $(x, y, z) \rightarrow (x, y, 2z - 1)$ for the second.

Let $[t_1, f_1, g]$ and $[t_2, f_2, g]$ be two tangles with flat connection in their complements such that $\partial_+ t_1 = \partial_- t_2$ and $f_1|_{\pi_1(\mathbb{R}^2 \setminus \partial_+ t_1, p(x_0))} = f_2|_{\pi_1(\mathbb{R}^2 \setminus \partial_- t_2, p(x_0))}$ and f_1 and f_2 have the same ramifications. Define the mapping $f_{1 \circ 2}$

$$f_{1 \circ 2}|_{\mathbb{R}^2 \times [0, 1/2]} = \tilde{f}_1, \quad f_{1 \circ 2}|_{\mathbb{R}^2 \times [1/2, 1]} = \tilde{f}_2,$$

where we identified elements of fundamental groups using (4) and (5). The map $f_{1 \circ 2}$ is a representation of the fundamental group of the complement of $t_1 \circ t_2$. We define the composition of two tangles with flat connections $[t_1, f_1, g]$ and $[t_2, f_2, g]$ as the equivalence class $[t_1 \circ t_2, f_{1 \circ 2}, g]$ and denote it as $[t_1, f_1, g] \circ [t_2, f_2, g]$. It is clear that the composition of two tangles with flat connections defined this way does not depend on the choice of representatives in $[t_1, f_1, g]$, $[t_2, f_2, g]$, and it is associative.

3.3 Tensor product

Let $[t_1]$ and $[t_2]$ be isotopy classes of two standard tangles. Let $|X|$ be the cardinality of a finite set X . Consider the map $\varphi : \mathbb{R}^2 \times I \rightarrow \mathbb{R}^2 \times I$ acting at the boundary as

$$(x, y, 0) \mapsto (x, y + |\partial_- t_1|, 0)$$




Figure 7:

$$(x, y, 1) \mapsto (x, y + |\partial_+ t_1|, 1)$$

Let $x_0(i)$ and $\tilde{x}_0(i)$ be the reference points for t_i ($i = 1, 2$). Assume that the map φ brings a representative t_2 of $[t_2]$ to t'_2 and that it brings $x_0(2)$ to $\tilde{x}_0(1)$. Choose representatives of t_1 and t_2 in such a way that one dimensional submanifolds t_1 and t'_2 do not tangle with each other (that their diagrams do not intersect). The isotopy class of $t_1 \cup t'_2$ is called the tensor product $[t_1] \otimes [t_2]$ of $[t_1]$ and $[t_2]$ [T-1]. It is clear that $[t_1] \otimes [t_2]$ does not depend on the choice of representatives.

Let $[t_1, f_1]$ and $[t_2, f_2]$ be two tangles with given representations of the fundamental group of the complement. Let t_1 and t'_2 be geometrical tangles as above. It is clear that $\pi_1(\mathbb{R}^2 \times I \setminus t_1 \cup t'_2, x_0(1))$ is isomorphic to the group freely generated by $\pi_1(\mathbb{R}^2 \times I \setminus t_1, x_0(1))$ and $\pi_1(\mathbb{R}^2 \times I \setminus t'_2, x_0(2))$. Definition of the tensor product will depend on how we embed groups $\pi_1(\mathbb{R}^2 \times I \setminus t_i, x_0(i))$ into $\pi_1(\mathbb{R}^2 \times I \setminus t_1 \cup t'_2, x_0(1))$.

Fix an isomorphism by identifying elements of $\pi_1(\mathbb{R}^2 \times I \setminus t_1 \cup t'_2, x_0(1))$ with corresponding elements of $\pi_1(\mathbb{R}^2 \times I \setminus t_1, x_0(1))$ and $\pi_1(\mathbb{R}^2 \times I \setminus t'_2, x_0(2))$ in accordance with Fig. 7. This identification means that we identify elements of $\pi_1(\mathbb{R}^2 \times I \setminus t'_2, x_0(2))$ with elements of $\pi_1(\mathbb{R}^2 \times I \setminus t_1 \cup t'_2, x_0(1))$ by gluing them with the ramification path for t_1 . For a pair of group ramified homomorphisms $f_i : \pi_1(\mathbb{R}^2 \times I \setminus t_i, x_0) \rightarrow G$, $i = 1, 2$, with ramifications g_1 and g_2 respectively define the group homomorphism

$$f_{1 \otimes_2} : \pi_1(\mathbb{R}^2 \times I \setminus t_1 \cup t'_2, x_0) \rightarrow G$$

by assigning

$$f_{1 \otimes_2}(\gamma) = \begin{cases} f_1(\gamma) & \text{if } \gamma \in \pi_1(\mathbb{R}^2 \times I \setminus t_1, x_0) \\ g_1 f_2(\gamma) g_1^{-1} & \text{if } \gamma \in \pi_1(\mathbb{R}^2 \times I \setminus t'_2, x'_0) \end{cases}$$

The ramification of $f_{1 \otimes_2}$ we will define as $g_1 g_2 \in G$. The equivalence class of $(t_1 \cup t'_2, f_{1 \otimes_2})$ will be called the tensor product of $[t_1, f_1, g_1]$ and $[t_2, f_2, g_2]$ and will be denoted $[t_1, f_1, g_1] \otimes [t_2, f_2, g_2]$. This operation is associative.




Figure 8:

3.4 Framing

A framing of a geometric tangle t is a continuous nonsingular section of the normal bundle to t . A framing of a tangle is the homotopy class of a geometric framing. We assume that the framing is parallel to the x -axis at $\partial_{\pm}t$, has positive projection to this axis and unit length.

3.5 The category of framed G -tangles

Here G -framed tangle is a tangle with a ramified representation of the fundamental group of the complement.

Define the category of framed G -tangles $\widetilde{\mathcal{FT}}(G)$ as follows:

- *Objects* are pairs $(\emptyset; f)$ where $f : \pi_1(\mathbb{R}^2) \rightarrow e \in G$ with the trivial ramification ($g = 1$) and $((\epsilon_1, \dots, \epsilon_n); f, g)$ where $\epsilon_i = \pm 1$, f is a representation of the fundamental group of $\mathbb{R}^2 \setminus \{(0, 1), \dots, (0, n)\}$ with the ramification g .
- *Morphisms* between $((\epsilon_1, \dots, \epsilon_n); f_+; g_+)$ and $((\sigma_1, \dots, \sigma_k); f_-; g_-)$ are nontrivial only if $g_+ = g_-$. They are oriented framed tangles with flat connections (with ramified representation of the fundamental group of the complement) $[t, f, g]$ where t is a tangle with $|\partial_- t| = k$, $|\partial_+ t| = n$ and f is a flat connection in the complement of t such that $f|_{\pi_1(\mathbb{R}^2 \setminus \partial_{\pm} t, x_0)} = f_{\pm}$. The ramification g is defined by the ramifications of the objects $g = g_+ = g_-$. The orientation of boundary components should agree with signs ϵ_i, σ_j as is shown on Fig. 8.
- The composition is defined above in Subsection 3.2.
- The identity element in $\text{End}((\epsilon_1, \dots, \epsilon_n), f, g)$ consists of the identity tangle (see Fig. 9) with the representation of the fundamental group and the ramification uniquely defined by its values on the boundary.

The category $\widetilde{\mathcal{FT}}(G)$ is braided monoidal with the tensor product described above.




Figure 9:




Figure 10:

The braiding is given by the tangle corresponding to the diagram from Fig. 10. The corresponding ramified representation of the fundamental group of its complement is completely determined by the boundary values which are fixed by the choice of objects.

3.6 The combinatorial description of $\widetilde{\mathcal{FT}}(G)$

The category $\widetilde{\mathcal{FT}}(G)$ can be described in terms of equivalence classes of diagrams with elements of G assigned to edges of the diagram.

Representations of the fundamental group of the complement to a tangle are in bijections with maps $E(D_t) \rightarrow G$ which satisfy Wirtinger relations. Therefore a tangle with a flat connection in the complement can be regarded as an equivalence class of diagrams equipped with such data modulo Reidemeister moves. If the tangle is framed, one has to consider only framed Reidemeister moves (see Figures 11, 12 13).

A representation of the fundamental group of the complement to n distinct points in $\mathbb{R}^2$ is fixed uniquely by the monodromies along a system of fundamental cycles. Choose them as on Fig. 14 then an object of the category $\widetilde{\mathcal{FT}}(G)$ can be written as a collection $(\epsilon_1, \dots, \epsilon_n; g_1, \dots, g_n; g)$ where $g_i \in G$ is a monodromy along the cycle shown on Fig. 14 around the point $(0, i, 0)$ and g is the ramification.

Thus, we can describe the category $\widetilde{\mathcal{FT}}(G)$ as follows:




Figure 11:




Figure 12:

- *Objects* are collections $(\epsilon_1, \dots, \epsilon_n; g_1, \dots, g_n; g)$, where $\epsilon_i = \pm$ and $g_i, g \in G$.
- *Morphisms* between the object $(\epsilon_1, \dots, \epsilon_n; g_1, \dots, g_n; g)$ and the object $(\sigma_1, \dots, \sigma_m; h_1, \dots, h_m; g)$ are diagrams of oriented framed tangles with elements of G assigned to edges modulo Reidemeister moves and Wirtinger relations. The orientation and group elements assigned to boundary components should agree the orientation and group elements of corresponding boundary components.
- *Composition* is the usual composition of diagrams of tangles with group elements assigned to edges.
- *Tensor product* for objects is $(\{\epsilon_i\}; \{g_i\}; g) \otimes (\{\sigma_i\}; \{h_i\}; h) = (\{\epsilon_i\}\{\sigma_i\}; \{g_i\}\{h_i\}; gh)$. The tensor product of two morphisms $(D_1, \{g_e\}, g)$ and $(D_2, \{h_e\}, h)$ is the diagram which $D_1 \sqcup D_2$ is the tensor product of two diagrams (juxtaposition) with the elements g_e assigned to the edges of D_1 and elements $g^{-1}h_e g$ assigned to edges of D_2 . The ramification of the tensor product is gh .

3.7 Combinatorial description of $\widetilde{\mathcal{FT}}(G)$ for factorizable G

Let G be a factorizable group. According to Theorem 1 a G -coloring of diagram D_t is in one to one correspondence with a flat G -connection in the complement to tangle t by assigning




Figure 13:




Figure 14:

monodromies γ_e to edges of the diagram:

$$\{x_e\}_{e \in D_t} \mapsto \{\gamma_e = (x_{e_1})_-^{\epsilon_1} \dots (x_{e_n})_-^{\epsilon_n} x_e (x_{e_n})_-^{-\epsilon_n} \dots (x_{e_1})_-^{-\epsilon_1}\}_{e \in D_t} \quad (6)$$

More precisely, it is a bijection for factorizable groups and a bijection on an open dense subset of $G^{\times E(D_t)}$ for almost factorizable Lie groups. This suggests the description of the the category $\widetilde{\mathcal{FT}}(G)$ in term of G -colored diagrams.

Define the category $\mathcal{FT}(G)$ of G -colored framed diagrams as follows:

- *Objects* are sequences $\{(\epsilon_1, x_1), \dots, (\epsilon_n, x_n)\}$ where $\epsilon_i = \pm 1$ and generic $x_i \in G$.
- *Morphisms* are G -colored framed diagrams modulo framed Reidemeister moves (see Figures 11, 12, 13)

This category is monoidal, braided. The composition is obvious. The tensor product is essentially the same as the tensor product of diagrams of framed tangles (see for example [T-1]). The decoration of the tensor product is defined by the decoration of factors.

Let $\widetilde{\mathcal{FT}}$ be the category of tangles with flat connections in the complement with the standard ramification (see section 2.4).

Proposition 1. *Consider the map $E : \mathcal{FT}(G) \rightarrow \widetilde{\mathcal{FT}}(G)$ acting on objects as*

$$\{(\epsilon_1, x_1), \dots, (\epsilon_n, x_n)\} \rightarrow \{(\epsilon_1, g_1), \dots, (\epsilon_n, g_n)\}$$




Figure 15:

where

$$g_i = (x_1)_-^{\epsilon_1} \dots (x_{i-1})_-^{\epsilon_{i-1}} x_i (x_{i-1})_-^{-\epsilon_{i-1}} \dots (x_1)_-^{-\epsilon_1}$$

and on morphisms as $(D_t \text{ with a } G\text{-coloring}) \rightarrow (t \text{ with ramified representations of the } \pi_1 \text{ of the complement to } t \text{ defined by the } G\text{-coloring of } D_t)$. It is a monoidal, braided functor $\mathcal{FT}(G) \rightarrow \widetilde{\mathcal{FT}}(G)$. This functor is also an equivalence of categories.

The proof follows from the bijection between tangles with given representation of the fundamental group of the complement and the equivalence classes of G -colored diagrams.

Because the categories $\widetilde{\mathcal{FT}}(G)$ and $\mathcal{FT}(G)$ are naturally equivalent we will abuse notations and will denote both of them by $\mathcal{FT}(G)$.

When G is an almost factorizable Lie group the category of G -colored diagrams has the tensor product, but the dual object exists only on an open dense subset of objects. The braiding in this case also exists only on an open subset of objects of this category.

3.8 Elementary diagrams

The following fact is a key for construction of invariants of tangles via braided monoidal categories.

Proposition 2. *Framed G -colored tangles are compositions of tensor products of elementary diagrams. Elementary diagrams are given on Fig. 15 .*

The proof of this proposition and the definition of elementary diagrams are the same as for the category of framed tangles (see [T-1]).

4 Holonomy Yang-Baxter equation and invariants of tangles with flat connections

Let G be a factorizable group (or a factorizable Lie group).

4.1 Holonomy Yang-Baxter equation

Assume we have the following data:

- Collection of finite-dimensional vector spaces $\mathfrak{X}$
- For each pair (X, Y) of vector spaces from this collection we have a map

$$R^{X,Y} : G \times G \rightarrow \text{End}(X \otimes Y) \quad (7)$$

(or, more generally, a section of a vector bundle over $G \times G$ with the fiber $X \otimes Y$).

Let $x_L, x_R : G \times G \rightarrow G$ be mappings acting as

$$x_L(x, y) = x_- y x_-^{-1}, \quad x_R(x, y) = x_L(x, y)_+^{-1} x x_L(x, y)_+ \quad (8)$$

Definition 2. *The data described above define a system of holonomy R -matrices if the equation*

$$R_{12}^{X,Y}(x_R(x, x_L(y, z)), x_R(y, z)) R_{13}^{X,Z}(x, x_L(y, z)) R_{23}^{Y,Z}(y, z) = \quad (9)$$

$$R_{23}^{Y,Z}(x_L(x, y), x_L(x_R(x, y), z)) R_{13}^{X,Z}(x_R(x, y), z) R_{12}^{X,Y}(x, y) \quad (10)$$

holds for any generic triple $x, y, z \in G$ and any $X, Y, Z \in \mathfrak{X}$. Here the mappings $x_{L,R}$ are as above, all factors act in $X \otimes Y \otimes Z$ and lower indices in the R -matrices indicate in which factors of the tensor product they act nontrivially.

We will call the equation (9) the *holonomy Yang-Baxter equation*.

For a linear operator $f : V \rightarrow V$ we denote by f^t the dual linear operator $f^t : V^* \rightarrow V^*$.

Definition 3. *Let V and W be two finite-dimensional vector spaces and W^* be the vector space dual to W . We will say that the linear map $R : V \otimes W \rightarrow V \otimes W$ is a cross-nondegenerate if the corresponding linear map $R^{t^2} : V \otimes W^* \rightarrow V \otimes W^*$ is nondegenerate.*

For cross-nondegenerate holonomy R -matrix acting in $X \otimes X$ define the map (more generally a section of the trivial vector bundle) $d^X : G \rightarrow \text{End}(X)$.

$$d^X(a) = (\text{tr} \otimes \text{id})(P((R^{t_1}(a, i(a)^{-1})^{-1})^{t_1}))$$

Here and below we will use the notation t_i for taking dual to an operator acting in a tensor product with respect to the i -th factor.

Lemma 1. *For generic $a \in G$ the linear operator $d^X(a)$ is invertible and*

$$d^X(a)^{-1} = (\text{id} \otimes \text{tr})((((R(i(a)^{-1}, a)^{-1})^{t_2})^{-1})^{t_2} P)$$

Proposition 3. *For $X, Y \in \mathfrak{X}$ and generic $x, y \in G$ the following identities hold*

$$R_{12}^{X,Y}(x, y) = d_2(a)^{-1}(((R_{12}^{X,Y}(x, y)^{-1})^{t_2})^{-1})^{t_2} d_2(y) \quad (11)$$

$$R_{12}^{X,Y}(a, b) = d_1(y)^{-1}(((R_{12}^{X,Y}(a, b)^{t_1})^{-1})^{t_1})^{-1} d_1(a) \quad (12)$$

$$d_1^X(b) d_2^Y(a) R^{X,Y}(x, y) = R^{X,Y}(x, y) d_1^X(x) d_2^Y(y) \quad (13)$$

where $a = x_L(x, y)$ and $b = x_R(x, y)$.

For $X \in \mathfrak{X}$ and generic $x \in G$ define the linear operator $w_X(x) \in \text{End}(X)$ as

$$w_X(x) = \text{tr}_1 \text{tr}_2 (P_{12}((R_{12}^{-1}(x, b)^{t_2})^{-1})^{t_2} P_{13} R_{13}(b, x)) \quad (14)$$

where $b = i(x^{-1})$.

Proposition 4. *For each $X, Y \in \mathfrak{X}$ we have*

$$(1 \otimes w_Y(x_L(x, y))) R^{X,Y}(x, y) = R^{X,Y}(x, y) (1 \otimes w_Y(y))$$

$$(w_X(x_R(x, y)) \otimes 1) R^{X,Y}(x, y) = R^{X,Y}(x, y) (w_X(x) \otimes 1)$$

The proof of this proposition and of the lemma are very similar. Each of the equalities follows from a sequence of equalities where each one corresponds to an elementary moves of corresponding diagrams. They are completely parallel to the proofs of similar statements from [R-1] (see Fig. 8, Fig. 13-17 from that paper and corresponding sequences of equalities). The only difference is that G -coloring should be taken into account.

4.2 The category $\mathcal{FT}(G, \mathfrak{X})$

Let $(D, \{x_e\}_{e \in E(D_t)})$ be a G -colored diagram of a tangle. We will call it *decorated by vector spaces* $\mathfrak{X}$ if a vector space from $\mathfrak{X}$ is assigned to each connected component of the corresponding tangle.

Define the category $\mathcal{FC}(G, \mathfrak{X})$ of $\mathfrak{X}$ -decorated G -colored diagrams as follows.

- *Objects* of $\mathcal{FC}(G, \mathfrak{X})$ are sequences $\{(\varepsilon_1, g_1, X_1), \dots, (\varepsilon_n, g_n, X_n)\}$ where $\varepsilon_i = \pm 1$, $g_i \in G' \subset G$ (recall that $G' \subset G$ is the set of elements in G which admit left and right factorization) and $X_a \in \mathfrak{X}$.
- *Morphisms* between two objects $\{(\varepsilon_a, g_a, X_a)\}_{a=1}^n$ and $\{(\sigma_a, h_a, Y_a)\}_{a=1}^k$ are G -colored diagrams decorated by vector spaces from $\mathfrak{X}$ modulo framed Reidemeister moves. Decorations of boundary components are determined by the initial and the target object. Elements g_a and h_a are colorings of boundary edges of such diagrams.
- Composition of morphisms is defined by gluing decorated diagrams along the common boundary.

It is clear that Ad_G of the G -coloring of a morphism in this category can be naturally identified with a gauge class of a flat connections in the complement to the corresponding tangle with connected components colored by $\mathfrak{X}$. The category $\mathcal{FT}(G, \mathfrak{X})$ is braided monoidal. The tensor product and the braiding are essentially the same as in the category $\mathcal{FT}(G)$.

4.3 The functor $F : \mathcal{FT}(G, \mathfrak{X}) \rightarrow \underline{Vect}$

Here we will show how to use a system of holonomy R -matrices to construct invariants of tangles with flat G -connections in the complement. The construction is similar to the one of invariants of tangles given in [RT]. We will construct the monoidal functor from the category $\mathcal{FC}(G, \mathfrak{X})$ to the category of all vector spaces.

Theorem 2. *There exists a unique covariant functor $F : \mathcal{FT}(G, \mathfrak{X}) \rightarrow \underline{Vect}$ such that*

- $F(\{(\varepsilon_1, g_1, X_1), \dots, (\varepsilon_n, g_n, X_n)\}) = X_1^{\varepsilon_1} \otimes \dots \otimes X_n^{\varepsilon_n}$, where $X^+ = X$, $X^- = X^*$.
- F is a monoidal functor, i.e.

$$F(\hat{D}_1 \otimes \hat{D}_2) = F(\hat{D}_1) \otimes F(\hat{D}_2)$$

where $\hat{D}_i$ are decorated diagrams.

- Values of F on elementary diagrams are:

1. If $I^\varepsilon : (\varepsilon, X, x) \rightarrow (\varepsilon, X, x)$ is the identity morphism then

$$F(I^\varepsilon) = id_{X^\varepsilon}.$$

2. For the morphism $e^r : (\varepsilon, X, x)^* \otimes (\varepsilon, X, x) \rightarrow 1$ we have:

$$F(e^r) = \begin{cases} e_X : X^* \otimes X \rightarrow \mathbb{C}, l \otimes v \mapsto l(d_X(x)v), & \text{if } \varepsilon = +1 \\ X \otimes X^* \rightarrow \mathbb{C}, v \otimes l \mapsto l(v), & \text{if } \varepsilon = -1 \end{cases}$$

3. For $i^r : 1 \rightarrow (\varepsilon, X, x) \otimes (\varepsilon, X, x)^*$ we have

$$F(i^r) = \begin{cases} i_X : \mathbb{C} \rightarrow X \otimes X^*, 1 \mapsto \sum_i d_X(x)^{-1} e_i \otimes e^i, & \text{if } \varepsilon = +1 \\ \mathbb{C} \rightarrow X^* \otimes X, 1 \mapsto \sum_i e^i \otimes e_i, & \text{if } \varepsilon = -1 \end{cases}$$

Here e_i is a linear basis in X and e^i is the dual linear basis in X^* .

4. If $c : (+, X, x) \otimes (+, Y, y) \rightarrow (+, Y, y) \otimes (+, X, x)$ is the braiding morphism,

$$F(c) = P^{X,Y} R^{X,Y}(x, y) : X \otimes Y \rightarrow Y \otimes X.$$

$$F(c^{-1}) = (P^{X,Y} R^{X,Y}(x, y))^{-1} : X \otimes Y \rightarrow Y \otimes X.$$

Here $P^{V,W} : V \otimes W \rightarrow W \otimes V$ is the linear map $P^{V,W}(x \otimes y) = y \otimes x$.

The proof of this theorem is completely parallel to the proof from of the corresponding theorem for tangles without flat connections (see [RT]). One should check the defining relations between elementary tangles, which is a routine exercise.

The value of F on a diagram is a linear map between the vector spaces which are defined by the decoration of the boundary of the diagram. The value of the functor F on a diagram depends only on its isotopy class and therefore F is an invariant of framed tangles. In the next section we will show that the functor F is gauge invariant.

5 Gauge invariance

5.1 The gauge group action and the dressing action

Let $g_1, \dots, g_n$ be a G -coloring of edges $e_1, \dots, e_n$ on Fig. 16 and let γ_i be the holonomy along the path Γ_i . For generic colorings we have:

$$\gamma_i = (g_1)_+^{\epsilon_1} \dots (g_i)_+^{\epsilon_i} (g_i)_-^{-\epsilon_i} \dots (g_1)_-^{-\epsilon_1} \quad (15)$$




Figure 16:

or, equivalently,

$$\epsilon_i(g_i) = (\gamma_{i-1})_+^{-1} \gamma_i (\gamma_{i-1})_- \quad (16)$$

Here ϵ_i is the orientation of the edge e_i (+ for going up and $-$ for going down) and

$$\epsilon(g) = \begin{cases} g & \text{for } \epsilon = +1 \\ i(g) & \text{for } \epsilon = -1 \end{cases}$$

Holonomies γ_i and colorings g_i can be regarded as an element of $G^{\times n}$. Then formulae (15), (16) describe two mutually inverse birational mappings of $G^{\times n}$ to itself.

The gauge group acts on γ_i by diagonal conjugation in $G^{\times n}$:

$$x : \gamma_i \mapsto x \gamma_i x^{-1} \quad (17)$$

The gauge action of G on this space is the diagonal action by conjugations.

For $x \in G$ consider the (birational) mapping $\alpha_x : G^{\times n} \rightarrow G^{\times n}$ defined as:

$$\epsilon_i(\alpha_x(g_i)) = x^{\gamma_{i-1}} \epsilon_i(g_i) (x^{\gamma_{i-1}})^{-1} \quad (18)$$

where γ_i are as above and

$$x^\gamma = (x \gamma x^{-1})_+^{-1} x \gamma_+ = (x \gamma x^{-1})_-^{-1} x \gamma_- \quad (19)$$

is the *dressing* action of γ on x .

Notice that the dressing action preserves the subgroups $G_\pm$ and

$$(x_+)^{\gamma} = (\gamma_-^{-1} x_+^{-1})_+^{-1} \quad (20)$$

$$(x_-)^{\gamma} = (x_- \gamma_+)_-^{-1} \quad (21)$$

Proposition 5. *The mapping (16) intertwines the gauge action of $x \in G$ and the mapping α_x .*




Figure 17:

Proof. Let $g^x = \{(g^x)_1, \dots, (g^x)_n\}$ be the result of the gauge action of $x \in G$ on g -coordinates. We have:

$$\begin{aligned} \epsilon_i((g^x)_i) &= (x\gamma_{i-1}x^{-1})_+^{-1}x\gamma_ix^{-1}(x\gamma_{i-1}x^{-1})_- \\ &= (x\gamma_{i-1}x^{-1})_+^{-1}x(\gamma_{i-1})_+\epsilon_i(g_i)(\gamma_{i-1})_-^{-1}x^{-1}(x\gamma_{i-1}x^{-1})_- = x^{\gamma_{i-1}}\epsilon_i(g_i)(x^{\gamma_{i-1}})^{-1} = \epsilon_i(\alpha_x(g)_i). \end{aligned}$$

Here we used formulae (19) for the dressing action. $\square$

Corollary 1. *If x belongs to one of the subgroups $G_{\pm} \subset G$ we have*

$$\epsilon_i(\alpha_{x_+}(g)_i) = x_+^{(g_{i-1})_-^{-\epsilon_{i-1}} \dots (g_1)_-^{-\epsilon_1}} \epsilon_i(g_i) \left(x_+^{(g_{i-1})_-^{-\epsilon_{i-1}} \dots (g_1)_-^{-\epsilon_1}} \right)^{-1} \quad (22)$$

$$\epsilon_i(\alpha_{x_-}(g)_i) = x_-^{(g_1)_+^{\epsilon_1} \dots (g_{i-1})_+^{\epsilon_{i-1}}} \epsilon_i(g_i) \left(x_-^{(g_1)_+^{\epsilon_1} \dots (g_{i-1})_+^{\epsilon_{i-1}}} \right)^{-1} \quad (23)$$

Lemma 2. *Let x, y, a, b be G -colorings of the diagram from Fig. 17. Then,*

$$\epsilon_e(a) = x_+ \epsilon_e(y) x_+^{-1}, \quad b_+ = (y_-^{-\epsilon_e} x_+^{-1})_+^{-1}$$

Proof. According to the definition of G -coloring we have:

$$b = a_-^{-\epsilon_e} x a_-^{\epsilon_e}, \quad \epsilon_e(a) = x_+ \epsilon_e(y) x_+^{-1}$$

This gives the formula for a . In particular,

$$a_- = (y_-^{-\epsilon_e} x_+^{-1})_-^{\epsilon_e}$$

so that

$$b_+ = (a_-^{-\epsilon_e} x_+)_+ = \left((y_-^{-\epsilon_e} x_+^{-1})_-^{-1} x_+ \right)_+ = \left((y_-^{-\epsilon_e} x_+^{-1})_+^{-1} y_-^{-\epsilon_e} x_+^{-1} x_+ \right)_+ = (y_-^{-\epsilon_e} x_+^{-1})_+^{-1}$$

$\square$




Figure 18:

Lemma 3. *Let x, y, a, b be colorings the diagram from Fig. 18. Then*

$$\epsilon_e(a) = x_-^{-1} \epsilon_e(y) x_-, \quad b_- = (x_-^{-1} y_+^{\epsilon_e})_-$$

Proof. From the definition of G -coloring we have

$$\epsilon_e(a) = x_-^{-1} \epsilon_e(y) x_-, \quad b = y_+^{-\epsilon_e} x y_+^{\epsilon_e}$$

so the required formulas are straightforward. □

Proposition 6. *If $g_1, \dots, g_n$ are G -colorings of "lower" edges of diagrams in Figures 19, 20, then the colorings of "upper" edges are*

$$\epsilon_i(\tilde{g}_i) = x_+^{(g_{i-1})_-^{-\epsilon_{i-1}} \dots (g_1)_-^{-\epsilon_1}} \epsilon_i(g_i) \left(x_+^{(g_{i-1})_-^{-\epsilon_{i-1}} \dots (g_1)_-^{-\epsilon_1}} \right)^{-1}$$

for the diagram in Fig 19 and

$$\epsilon_i(\tilde{g}_i) = (x_-^{-1})^{(g_1)_+^{\epsilon_1} \dots (g_{i-1})_+^{\epsilon_{i-1}}} \epsilon_i(g_i) \left((x_-^{-1})^{(g_1)_+^{\epsilon_1} \dots (g_{i-1})_+^{\epsilon_{i-1}}} \right)^{-1}$$

for the diagram in Fig 20.

Proof. Let us consider first the diagram from Fig.19. Lemma 2 implies the following relations between G -colorings:

$$\epsilon_i(\tilde{g}_i) = (x_{i-1})_+ \epsilon_i(g_i) (x_{i-1})_+^{-1}, \quad (x_i)_+ = ((g_i)_-^{-\epsilon_i} (x_{i-1})_+^{-1})_+^{-1}$$

From here we deduce that

$$(x_i)_+ = ((g_i)_-^{-\epsilon_i} \dots (g_1)_-^{-\epsilon_1} x_+^{-1})_+^{-1} = x_+^{(g_i)_-^{-\epsilon_i} \dots (g_1)_-^{-\epsilon_1}}$$




Figure 19:




Figure 20:

This, together with the formula (20) imply the first statement of the proposition.

Similarly, for the G -coloring of the diagram from Fig. 20 we have

$$\epsilon_i(\tilde{g}_i) = (x_{i-1})_-^{-1} \epsilon_i(g_i) (x_{i-1})_-, \quad (x_i)_- = ((x_{i-1})_-^{-1} (g_i)_+^{\epsilon_i})_-$$

This gives the formula for $(x_i)_-$:

$$(x_i)_- = (x_-^{-1} (g_1)_+^{\epsilon_1} \dots (g_i)_+^{\epsilon_i})_- = \left((x_-^{-1})^{(g_1)_+^{\epsilon_1} \dots (g_i)_+^{\epsilon_i}} \right)^{-1}$$

Together with the formula (21) this implies the last statement of the proposition and completes the proof. □

5.2 Gauge invariance of the functor F

Let $R^{X,Y}(x, y)$ be a holonomy R -matrix acting in $X \otimes Y$ where $X, Y \in \mathfrak{X}$. Define the operator

$$\sigma(R^{X,Y}(x, y)) = P^{YX} R^{X,Y}(x, y) P^{XY} : Y \otimes X \rightarrow X \otimes Y$$

Let t be an $\mathfrak{X}$ -decorated G -colored tangle. Assume that

$$t \in \text{Hom}_{\widetilde{\mathcal{FT}}(G, \mathfrak{X})}((\epsilon_1, Y_1, y_1) \otimes \dots \otimes (\epsilon_n, Y_n, y_n), (\sigma_1, Z_1, z_1) \otimes \dots \otimes (\sigma_k, Z_k, z_k))$$

Denote by $F(t)$ the value of functor F described in Subsection 4.3.

Theorem 3. *Let t^x be an $\mathfrak{X}$ -decorated G -colored tangle with the G -coloring obtained by the gauge action of $x \in G$ on the G -coloring of the tangle t . Then the following relations hold for $x_{\pm} \in G_{\pm}$*

$$(1 \otimes F(t^{x_+}))R_{0n}^{\varepsilon_n, n}(\{y\}, x)^{-1} \dots R_{01}^{\varepsilon_1, 1}(\{y\}, x)^{-1} = R_{0k}^{\sigma_k, k}(\{z\}, x)^{-1} \dots R_{01}^{\sigma_1, 1}(\{z\}, x)^{-1}(1 \otimes F(t)) \quad (24)$$

$$(F(t^{x_-}) \otimes 1)\tilde{R}_{1, n+1}^{\varepsilon_1, 1}(\{y\}, x)^{-1} \dots \tilde{R}_{n, n+1}^{\varepsilon_n, n}(\{y\}, x)^{-1} = R_{1, k+1}^{\sigma_1, 1}(\{z\}, x)^{-1} \dots R_{k, k+1}^{\sigma_k, k}(\{z\}, x)^{-1}(F(t) \otimes 1) \quad (25)$$

Both sides in (24) are linear maps $X \otimes Y_1 \otimes \dots \otimes Y_n \rightarrow X \otimes Z_1 \otimes \dots \otimes Z_k$ with

$$R_{0i}^{(+, i)}(\{y\}, x) = \sigma(R^{Y_i, X}(\alpha_{x_-}(y)_i, x_i))_{0i}$$

$$R_{0i}^{(-, i)}(\{y\}, x) = \sigma((R^{Y_i, X}(\alpha_{x_-}(y)_i, x_{i-1})^{t_1})^{-1})_{0i}$$

Here we used the notation

$$(a \otimes b)_{0i} = a_0 \otimes 1 \otimes \dots \otimes b_i \otimes \dots \otimes 1 \in \text{End}(X \otimes Y_1 \otimes \dots \otimes Y_n)$$

Both sides in (25) are linear maps $Y_1 \otimes \dots \otimes Y_n \otimes X \rightarrow Z_1 \otimes \dots \otimes Z_k \otimes X$ with

$$\tilde{R}_{i, n+1}^{(+, i)}(\{y\}, x) = \sigma(R^{X, Y_i}(x_{i-1}, \alpha_{x_+}(y)_i))_{i, n+1}$$

$$\tilde{R}_{i, n+1}^{(-, i)}(\{y\}, x) = \sigma((R^{X, Y_i}(x_i, \alpha_{x_+}(y)_i)^{-1})^{t_2})_{i, n+1}$$

where

$$(a \otimes b)_{i, n+1} = 1 \otimes \dots \otimes a_i \otimes \dots \otimes 1 \otimes b_{n+1}$$

The maps $\alpha_{x_{\pm}}$ are defined in (22) (23), and the elements $x_i, \tilde{x}_i, x^i, \tilde{x}^i$ are determined by the recurrence relations

$$x_{i-1} = (\alpha_{x_-}(y)_i)^{-\epsilon_i} x_i (\alpha_{x_-}(y)_i)^{\epsilon_i}$$

$$x^{i-1} = (y_i)_{+}^{\epsilon_i} x_i (y_i)_{+}^{-\epsilon_i}$$

with $x_0 = x_+$ and $x^0 = x_-$. Replacing y by z and n by k we will have the definition of the remaining R -matrices.




Figure 21:




Figure 22:

Proof. First let us prove the equation (24). Consider the diagram which is represented on the Fig. 22. The value of the functor F on this diagram coincide with the value of this functor on the diagram which obtained from it by moving the component colored by x under the tangle to top of the diagram. This equality is exactly the equality (24). Similarly, moving the component colored by x on the diagram from Fig. 21 we prove the second equality. Evaluation of the functor F on these diagrams is an easy routine. Formulae from Appendix 7.2 are useful.

Q.E.D.

Thus, the functor F is an invariant of tangles with flat G -connection in the complement which is transforms as it is described above with respect to the action of the gauge group.

Corollary 2. *If t is a link, the invariant $F(t)$ is a scalar, i.e. it takes values in the base field which was assumed $\mathbb{C}$. It is invariant with respect to the gauge group action on the flat connection and therefore is a function (or a section of a line bundle) on the moduli space of flat G -connections on the complement to the tangle t .*

6 Conclusion

We described a construction of invariants of tangles with flat connections in the complement which is based on a system of nondegenerate and cross-nondegenerate holonomy R -matrices. In the second paper of this series we will show how to construct holonomy R -matrices from the representation theory of quantum groups at roots of 1. We will also show how to use

these holonomy R -matrices to construct invariants of 3-manifolds with flat G -connection in it.

References

- [J] V. Jones A new knot polynomial and von Neumann algebras. Bull.Amer.Math.Soc.(N.S.)12(1985), no. 1, 103-111.
- [J-1] V. Jones Notes on a talk at Atiyah seminar (Nov. 1986).
- [R] N. Reshetikhin. Quantized universal enveloping algebras, Yang-Baxter equation and invariants of links. I and II.LOMI-preprints, E-87, E-17-87, 1988.
- [R-1] N. Reshetikhin. Quasitriangular Hopf algebras and invariants of links. *Leningrad Math. J.* 1 (1990), no. 2, 491–513
- [RT] N. Reshetikhin,V. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.* 127 (1990), no. 1, 1–26
- [T] V. Turaev The Yang-Baxter equation and invariants of links. *Inv.Math.* 92(1988), no. 3, 527-553.
- [T-1] V. Turaev. *Quantum invariants of knots and 3-manifolds.* de Gruyter Studies in Mathematics, 18. Walter de Gruyter & Co., Berlin, 1994.
- [T-2] V. Turaev. Homotopy field theory in dimension 3 and crossed group-categories. preprint GT/0005291.
- [XW] A. Weinstein and P. Xu Classical solutions of the quantum Yang-Baxter equation. *Comm. Math. Phys.* 148 (1992), no. 2, 309–343.

7 Appendix

7.1

Let us show that the crossing rules from Fig. 6 imply the rules in the definition of the G -coloring. Indeed, consider pairs of pathes from Fig. 23-26. Isotopy equivalence of these pathes imply the equality of corresponding monodromies. Computing monodromies along these pathes according to crossing rule from Fig. 6 we obtain the definition of the G -coloring.




Figure 23:




Figure 24:




Figure 25:




Figure 26:




Figure 27:




Figure 28:

7.2

Let t_1, t_2 and t_1^{-1}, t_2^{-1} be diagrams from Fig. 27-30. The functor F can be evaluated on them using the second framed Reidemeister. As a result we have the following linear maps:

$$F(t_1) = ((R^{VW}(b, y)^{-1})^{t_2})^{-1} P^{W^*V} : W^* \otimes V \rightarrow V \otimes W^*$$

$$F(t_2) = P^{W^*V} ((R^{WV}(a, x))^{t_1})^{-1} : W^* \otimes V \rightarrow V \otimes W^*$$

$$F(t_1^{-1}) = F(t_1)^{-1} = P^{VW^*} (R^{VW}(b, y)^{-1})^{t_2}$$

$$F(t_2^{-1}) = F(t_2)^{-1} = (R^{WV}(a, x))^{t_1} P^{VW^*}$$




Figure 29:




Figure 30: